

A Supplement to “On Normality of a Family of Holomorphic Functions”

Chikara WATANABE

*Department of Mathematics, College of Liberal Arts,
Kanazawa University.*

(Received May 29, 1974)

In our previous paper¹⁾, we proved the following :

Let D be a domain in C^n and let $\{f_j\}$ be a sequence of holomorphic functions in D such that

(i) ; $\{f_j\}$ is bounded at each point of D , i. e., $\{f_j(p)\}$ is a bounded subset of the complex plane C for each point $p \in D$,

(ii) ; the sequence $\{G_j\}$ of graphs of f_j converges analytically to an analytic set A in $D \times C$.

Then the sequence $\{f_j\}$ converges uniformly to a holomorphic function in D on every compact subset of D .

In this note we show that the condition (ii) cited above is weakened, that is we show the following

THEOREM. *Let D be a domain in C^n and let $\{f_j\}$ be a sequence of holomorphic functions in D such that*

(i) ; $\{f_j\}$ is bounded at each point of D ,

(ii) ; the sequence $\{G_j\}$ of graphs of f_j converges geometrically to a proper analytic set A in $D \times C$.

Then $\{f_j\}$ converges uniformly to a holomorphic function in D on every compact subset of D .

Proof. Let E be a set of non fine point of A , that is $p \notin E$ if and only if the set $A(p) = A \cap \{(p, w) \in C^{n+1}\}$ has no finite accumulating point. Then E is a proper analytic set in D ²⁾. Let $P_0 \in D - E$, then since $\{f_j(p_0)\}$ is bounded there exists at least one limit point q_0 . By the definition of the geometric convergence, it holds that $(p_0, q_0) \in A$. Since A is proper and since $p_0 \notin E$ we can take an open polydisc $\Delta \subset D - E$ with center at p_0 and an open disc $U \subset C$ with center at q_0 such that $A \cap (\Delta \times \partial U) = \emptyset$. Then there

1) On normality of a family of holomorphic functions, Publications RIMS, Vol 9, No. 3, 1974.

2) See *ibid.*, section 2.

exists a positive integer j_0 such that $G_j \cap (\Delta \times \partial U) = \emptyset$ and $G_j \cap (\Delta \times U) \neq \emptyset$ for all $j \geq j_0$. Let $\pi: \Delta \times U \rightarrow \Delta$ be the natural projection and put

$$\pi_j = \pi |_{G_j \cap (\Delta \times U)}: G_j \cap (\Delta \times U) \longrightarrow \Delta.$$

Then it is easily seen that π_j is a proper mapping, so that $(G_j \cap (\Delta \times U), \pi_j, \Delta)$ is an analytic cover³⁾. Thus π_j is onto. This means that $\{f_j\}$ is uniformly bounded on Δ , that is $\{f_j\}$ is locally uniformly bounded on $D - E$. Let $p_0 \in E$. Since E is a proper analytic set in D , by a linear change of coordinate if necessary, we may assume that there exists a polydisc Δ with center at p_0 such that E does not meet with the distinguished boundary of Δ . Then by the maximum principle of the holomorphic functions, $\{f_j\}$ is bounded on Δ . That is $\{f_j\}$ is locally uniformly bounded on D , and then $\{f_j\}$ is a normal family. If a subsequence $\{f_{\nu_j}\}$ of $\{f_j\}$ converges to a holomorphic function f on every compact set in D , then $\{G_{\nu_j}\}$ converges geometrically to the graph of f and $A = \text{graph of } f$. From this, $\{f_j\}$ itself converges uniformly to f on every compact set in D .

REMARK 1. Under the condition of the above theorem, the set E is in fact empty since A is a graph of a holomorphic function.

REMARK 2. The conditions (i), (ii) are said in other words as follows.

- (i); $\{f_j\}$ is bounded at each point of D and equicontinuous at some point $p_0 \in D$,
- (ii); the sequence $\{G_j\}$ converges geometrically to an analytic set A in $D \times C$.

In fact, we have only to show that A is proper. There exists an open polydisc $\Delta \subset D$ and a positive constant M such that $|f_j(p)| \leq M$ if $p \in \Delta$. Thus if $(p, q) \in A$ then $|q| \leq M$, that is A is a proper analytic set.

3) See, for example, Gunning R.C., and Rossi H., Analytic functions of several complex variables, Chapter III, B.